

# Ambulatory Use of Electronic Nicotine Delivery Systems - Redefining Topography Endpoints

Robert Underly, Evan Nudi, Tim Pionk, and Jeffrey Smith



#### **Current Topography Assessments**





#### In Clinic Use

- Confinement (with some combination of the below):
  - Fixed puff count
  - Fixed puff session length
  - Fixed time in clinic
  - Variable methods of quantification

#### **Natural Environment**

- Ambulatory Use
- "Sessions" of use
- Portable data capture device

# The Landscape of Topography Endpoints



Current Endpoints
Puff Duration
Puff Volume
Time Between Puffs
Flow Rate
Puffs Count

- Each endpoint is contingent on the test condition (natural environment vs confinement).
- Puffs counts are currently difficult to understand broadly as some studies hold this variable constant or limit subjects' time with the product.
- No study has currently reported cumulative data for endpoints over a day of use.

### **Product Use and Behavior (PUB) Instrument**





Parameter	Unit			
Puff Duration	Seconds (s) (battery active)			
Number of Puffs	Count (# of battery activations)			
Sessions of Use	Count			
Inter Puff Interval (IPI)	Seconds (s) (Latency between activations)			
Angle of Use	Puff initiation and conclusion (XYZ planes)			
Battery Characteristics	Voltage and Current			
with the addition of every data point collected is time/date stamped				



### **Product Use and Behavior (PUB) Instrument**




Product Characteristics with and without the PUB							
	Vuse Alto with PUB						
Product Battery Life (mean puffs)	209	210					
PUB Battery Life (mean puffs)	N/A	2000					
Idle Battery Life (hours)	>168	72					
Mean Puffs per Cartridge	304	310					
<b>Product Charge Time (minutes)</b>	55 ± 7	N/A					
PUB Charge Time (minutes)	N/A	15 - 60					
<b>Total Particulate Matter (55/30/3)</b>	6.46	6.09					
Pressure Drop (mmWg)	105	125					
Product Length (mm)	105	145					
Product Weight (grams)	22.6	42.8					
Activated Product Voltage	3.0	3.0					
Product Resistance	1.25	1.35					




#### **Product Use and Behavior Instrument**






#### **Cumulative Time-series Data**







#### **SEEQ** – Real-time Analysis








#### **Understanding the Stratification of Use**








Percentile	Mean Puffs per Day
0 - 10 <sup>th</sup>	N/A
10 <sup>th</sup> – 20 <sup>th</sup>	18.27
20 <sup>th</sup> - 30 <sup>th</sup>	34.89
$30^{th}-40^{th}$	57.92
40 <sup>th</sup> - 50 <sup>th</sup>	82.30
50 <sup>th</sup> - 60 <sup>th</sup>	104.29
60 <sup>th</sup> - 70 <sup>th</sup>	133.82
70 <sup>th</sup> - 80 <sup>th</sup>	162.23
80 <sup>th</sup> - 90 <sup>th</sup>	211.99
90 <sup>th</sup> - 100 <sup>th</sup>	305.04

Percentile Rank

#### **Understanding the Stratification of Use**







Percentile	Mean Puffs per Day
0 – 10 <sup>th</sup>	N/A
10 <sup>th</sup> – 20 <sup>th</sup>	18.27
$20^{th} - 30^{th}$	34.89
$30^{th}-40^{th}$	57.92
40 <sup>th</sup> - 50 <sup>th</sup>	82.30
50 <sup>th</sup> - 60 <sup>th</sup>	104.29
60 <sup>th</sup> - 70 <sup>th</sup>	133.82
$70^{th} - 80^{th}$	162.23
80 <sup>th</sup> - 90 <sup>th</sup>	211.99
90 <sup>th</sup> – 100 <sup>th</sup>	305.04

### **Defining Sessions of Use**




#### **Operational Definition for Session:**

A period of use in which consecutive puffs occur without exceeding the mean Inter-puff interval (IPI). Gaps between puffs that exceed the mean IPI are identified as Intersession Intervals.

An initial data "cleansing" step was performed in which the 95<sup>th</sup>-99<sup>th</sup> percentile of IPIs were removed (12 – 16 hour gaps in use) from each subject.

#### **Intersession Interval**





Intersession Interval						
Group	Mean (sec)	St. Dev.				
Low	177.09	± 451.52				
Mid	57.67	± 215.28				
High	22.09	± 99.68				

#### **Topography Endpoints**



Variable	Mean	St. Dev.
Mean Puffs Per Day	111.25	± 68.31
IPI (sec)	3671.66	± 6541.64
Mean Puff Duration (sec)	2.60	± 1.18

n=128

Using only the traditional topography endpoints for cumulative time-series data, the data provide limited insight into use behavior.

### **Topography Endpoints in the Context of Sessions**



Variable	Low	Low Users		Users	High Users	
variable	Mean	St. Dev.	Mean	St. Dev.	Mean	St. Dev.
Mean Daily Puffs	36.37	± 17.40	100.79	± 22.17	200.80	± 53.51
Mean Daily Sessions	6.83	± 6.51	22.05	± 13.09	58.75	± 31.29
Session Length (sec)	158.92	± 301.11	146.27	± 277.10	90.75	± 185.37
IPI with Sessions (sec)	51.00	± 72.22	45.67	± 48.68	35.06	± 38.03
Puffs within Sessions	5.02	± 7.30	4.44	± 6.08	3.40	± 4.30
Mean Puff Duration (sec)	2.36	± 1.43	2.80	± 1.27	2.49	± 1.07
Intersession Interval (min)	177.09	± 451.52	57.67	± 215.28	22.09	± 99.68

Low users: n=32; Mid Users: n=62; High Users: n=34

# **Topography Endpoints in the Context of Sessions**



Variable	Low	Low Users		Users	High Users	
Variable	Mean	St. Dev.	Mean	St. Dev.	Mean	St. Dev.
Mean Daily Puffs	36.37	± 17.40	100.79	± 22.17	200.80	± 53.51
Mean Daily Sessions	6.83	± 6.51	22.05	± 13.09	58.75	± 31.29
Session Length (sec)	158.92	± 301.11	146.27	± 277.10	90.75	± 185.37
IPI with Sessions (sec)	51.00	± 72.22	45.67	± 48.68	35.06	± 38.03
Puffs within Sessions	5.02	± 7.30	4.44	± 6.08	3.40	± 4.30
Mean Puff Duration (sec)	2.36	± 1.43	2.80	± 1.27	2.49	± 1.07
Intersession interval (min)	177.09	± 451.52	57.67	± 215.28	22.09	± 99.68

Low users: n=32; Mid Users: n=62; High Users: n=34

## **Topography Endpoints in the Context of Sessions**



Variable	Vuse	Solo	Vuse Ciro		Vuse Vibe		Vuse Alto	
variable	Mean	St. Dev.	Mean	St. Dev.	Mean	St. Dev.	Mean	St. Dev.
Mean Daily Puffs	184.41	± 97.99	205.05	± 161.02	247.66	± 154.49	200.80	± 53.51
Mean Daily Sessions	32.26	± 19.56	43.48	± 37.30	71.55	± 63.66	58.75	± 31.29
Session Length (sec)	156.98	± 313.12	99.46	± 222.86	60.33	± 144.29	90.75	± 185.37
IPI within Sessions (sec)	26.89	± 60.40	23.51	± 26.72	20.25	± 22.63	35.06	± 38.03
Puffs within Sessions	6.04	± 6.74	5.19	± 7.94	3.71	± 5.53	3.40	± 4.30
Mean Puff Duration (sec)	2.11	± 0.84	2.69	± 1.25	1.95	± 1.13	2.49	± 1.07
Intersession Interval (min)	57.66	± 260.97	33.55	± 157.17	20.59	± 121.70	22.09	± 99.68


Data from the 70<sup>th</sup> percentile of use. Solo: n=19; Ciro: n=16; Vibe: n=15; Alto: n=34

### **Advancing Topography**



- Ambulatory data collection, along with appropriate data analytics, allows for a more expansive scope of consumer use behavior.
- Understanding patterns of use may allow for the improvement of products to be more acceptable to smokers wanting to transition to products lower on the risk continuum
- Using these data to drive analytic testing (puff regimen) parameters can provide more accurate estimates of consumer exposure.

	Regimen	Volume	IPI	Duration	Rest Period
Current	Non- Intense	55	30	3	N/A
Cur	Intense	80	15	5	N/A
New	Non- Intense	55	30	3	60 sec*
	Intense	80	15	5	60 sec*



#### **Acknowledgements**

**Pionk** 





Nudi

Underly

**Smith** 

Original Patent Team

- Jeffrey Smith
- Steven Alderman
- Stephen Sears
- Rajesh Sur
- Sarah Baxter-Wright
- John Darnell
- Jason Hong
- Paul Nelson
- Elaine Round

- Clinical Team
  - Gary Dull
  - John Darnell
  - Erin Evans
  - Paul Nelson
  - Kristen Prevette
  - Peter Chen
  - Anzhi Kaszycki

- IM Team
  - Josh Cooke
  - Angela Mcveigh
  - Ken Little
- Carolina Medical Electronics
  - Bradley Brown